

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.


**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Amendments to the Claims

This listing of claims will replace all prior listings of claims in the application.

Listing of Claims

1. (Presently Amended) A method of preventing or treating inflammatory response associated with atherosclerosis or restenosis in a mammal, comprising administering to said mammal an effective amount of a compound selected from the group consisting of structures of Formula VI, Formula VII, Formula VIII and Formula IX, wherein Formula VI is:



or a pharmaceutically acceptable salt thereof wherein, A^{VI} is

- a) Cl,
- b) Br,
- c) CN, ss
- d) NO₂, or
- e) F;

R^{VI-1} is

- a) R^{VI-5} , or
- b) SO₂ R^{VI-9}

R^{VI-2} , R^{VI-3} and R^{VI-4} may be the same or different and are selected from the group consisting of:

- a) H,
- b) halo^{VI},
- c) aryl^{VI},
- d) S(O)_m R^{VI-6} ,
- e) (C=O) R^{VI-6} ,

- f) $(C=O) \cdot OR^{VI-9}$,
- g) cyano,
- h) het^{VI} , wherein said het^{VI} is bound via a carbon atom,
- i) OR^{VI-10} ,
- j) $Ohet^{VI}$,
- k) $NR^{VI-7}R^{VI-8}$
- l) SR^{VI-10} ,
- m) $Shet^{VI}$,
- n) $NHCOR^{VI-12}$,
- o) $NHSO_2R^{VI-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{VI-11} , OR^{VI-13} , SR^{VI-10} , SR^{VI-13} , $NR^{VI-7}R^{VI-8}$, halo, $(C=O)C_{1-7}$ alkyl, or SO_mR^{VI-9} , and
- q) R^{VI-3} together with R^{VI-2} or R^{VI-4} form a carbocyclic or VI -het which may be optionally substituted by $NR^{VI-7}R^{VI-8}$, or C_{1-7} alkyl which may be optionally substituted by OR^{VI-14} ;

R^{VI-5} is

- a) $(CH_2CH_2O)_iR^{VI-10}$,
- b) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $NR^{VI-7}R^{VI-8}$, R^{VI-11} , SO_mR^{VI-9} , or OC_{2-4} alkyl which may be further substituted by het^{VI} , OR^{VI-10} , or $NR^{VI-7}R^{VI-8}$, or
- c) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11} , $NR^{VI-7}R^{VI-8}$, $SO_m^{VI}R^{VI-9}$, or C_{1-7} alkyl optionally substituted by R^{VI-11} , $NR^{VI-7}R^{VI-8}$, or $SO_m^{VI}R^{VI-9}$;

R^{VI-6} is

- a) C_{1-7} alkyl,
- b) $NR^{VI-7}R^{VI-8}$,
- c) aryl VI , or
- d) het^{VI} , wherein said het^{VI} is bound via a carbon atom;

R^{VI-7} and R^{VI-8} are independently

- a) H,
- b) aryl^{VI},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of aryl^{VI}, $NR^{VI-10}R^{VI-10}$, R^{VI-11} , SO_mR^{VI-9} , $CONR^{VI-10}R^{VI-10}$, or halo, or;
- d) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11} , $NR^{VI-7}R^{VI-8}$, $SO_m^{VI}R^{VI-9}$, or C₁₋₇alkyl optionally substituted by R^{VI-11} , $NR^{VI-7}R^{VI-8}$, or $SO_m^{VI}R^{VI-9}$, or
- e) R^{VI-7} and R^{VI-8} together with the nitrogen to which they are attached form a het^{VI};

R^{VI-9} is

- a) aryl^{VI},
- b) het^{VI},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $NR^{VI-10}R^{VI-10}$, R^{VI-11} , SH, $CONR^{VI-10}R^{VI-10}$, or halo;

R^{VI-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VI-11} is

- a) OR^{VI-10} ,
- b) Ohet^{VI},
- c) Oaryl^{VI},
- d) CO_2R^{VI-10} ,
- e) het^{VI},
- f) ^{VI}-aryl^{VI},
- g) CN, or

- h) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11} , $NR^{VI-7}R^{VI-8}$, $SO_m^{IV}R^{VI-9}$, or C_{1-7} alkyl optionally substituted by R^{VI-11} , $NR^{VI-7}R^{VI-8}$, or SO_mR^{VI-9} ;

R^{VI-12} is

- a) H,
- b) het^{VI} ,
- c) $aryl^{VI}$,
- d) C_{3-8} cycloalkyl,
- e) methyl, or
- f) C_{2-7} alkyl optionally substituted by $NR^{VI-7}R^{VI-8}$ or R^{VI-11} ;

R^{VI-13} is

- a) $(P=O)(OR^{VI-14})_2$,
- b) $CO(CH_2)_n^{IV}CON(CH_3)-(CH_2)_nSO_3^-M^{VI+}$,
- c) an amino VI acid,
- d) $C(=O)aryl^{VI}$,
- e) $C(=O)C_{1-7}$ alkyl optionally substituted by $NR^{VI-7}R^{VI-8}$, $aryl^{VI}$, het^{VI} , CO_2H , or $O(CH_2)_nCO_2R^{VI-14}$, or
- f) $C(=O)NR^{VI-7}R^{VI-8}$

R^{VI-14} is

- a) H, or
- b) C_{1-7} alkyl;

each i^{VI} is independently 2, 3, or 4;

each n^{VI} is independently 1, 2, 3, 4 or 5;

each m^{VI} is independently 0, 1, or 2;

M^{VI} is sodium, potassium, or lithium;

$aryl^{VI}$ is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;


wherein any $aryl^{VI}$ is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO_2R^{VI-14} , CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl

which maybe further substituted by one to three $\text{SR}^{\text{VI-14}}$, $\text{NR}^{\text{VI-14}}\text{R}^{\text{VI-14}}$, $\text{OR}^{\text{VI-14}}$, or $\text{CO}_2\text{R}^{\text{VI-14}}$;

het^{VI} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VI-14}}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which maybe further substituted by one to three $\text{SR}^{\text{VI-14}}$, $\text{NR}^{\text{VI-14}}\text{R}^{\text{VI-14}}$, $\text{OR}^{\text{VI-14}}$, or $\text{CO}_2\text{R}^{\text{VI-14}}$;

wherein Formula VII is



VII

or a pharmaceutically acceptable salt thereof,

wherein

A^{VII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO_2 , or
- e) F;

$\text{R}^{\text{VII-1}}$ is

- a) aryl^{VII},
- b) $\text{S}(\text{O})_m^{\text{VII}}\text{R}^{\text{VII-6}}$,
- c) $(\text{C}=\text{O})\text{R}^{\text{VII-6}}$, with the proviso that if $\text{R}^{\text{VII-6}}$ is $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, then $\text{R}^{\text{VII-7}}$ and $\text{R}^{\text{VII-8}}$ do not both equal H,
- d) $(\text{C}=\text{O})\text{OR}^{\text{VII-9}}$,
- e) cyano,

- f) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- g) Ohet^{VII} ,
- h) $\text{NR}^{\text{VII}-7}\text{R}^{\text{VII}-8}$ with the proviso that $\text{R}^{\text{VII}-7}$ and $\text{R}^{\text{VII}-8}$ do not both equal H,
- i) $\text{SR}^{\text{VII}-10}$,
- j) Shet^{VII} ,
- k) $\text{NHCOR}^{\text{VII}-12}$,
- l) $\text{NHSO}_2\text{R}^{\text{VII}-12}$,
- m) $\text{C}_{1-7}\text{alkyl}$ which is partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VII}-11}$, $\text{OR}^{\text{VII}-13}$, $\text{SR}^{\text{VII}-10}$, $\text{SR}^{\text{VII}-13}$, $\text{NR}^{\text{VII}-7}\text{R}^{\text{VII}-8}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}\text{alkyl}$, or $\text{SO}_m\text{R}^{\text{VII}-9}$, or
- n) $\text{C}_{1-7}\text{alkyl}$ which is substituted by one or more substituents of the group $\text{R}^{\text{VII}-11}$, $\text{OR}^{\text{VII}-13}$, $\text{SR}^{\text{VII}-10}$, $\text{SR}^{\text{VII}-13}$, $\text{NR}^{\text{VII}-7}\text{R}^{\text{VII}-8}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}\text{alkyl}$, or $\text{SO}_m\text{R}^{\text{VII}-9}$;

$\text{R}^{\text{VII}-2}$ is

- a) H,
- b) halo,
- c) aryl^{VII} ,
- d) $\text{S}(\text{O})_m\text{R}^{\text{VII}-6}$,
- e) $(\text{C}=\text{O})\text{R}^{\text{VII}-6}$,
- f) $(\text{C}=\text{O})\text{OR}^{\text{VII}-9}$,
- g) cyano,
- h) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- i) $\text{OR}^{\text{VII}-10}$,
- j) Ohet^{VII} ,
- k) $\text{NR}^{\text{VII}-7}\text{R}^{\text{VII}-8}$,
- l) $\text{SR}^{\text{VII}-10}$,
- m) Shet^{VII} ,
- n) $\text{NHCOR}^{\text{VII}-12}$,
- o) $\text{NHSO}_2\text{R}^{\text{VII}-12}$, or
- p) $\text{C}_{1-7}\text{alkyl}$ which may be partially unsaturated and optionally substituted by one or more substituents

- of the group R^{VII-11} , OR^{VII-13} , SR^{VII-10} , SR^{VII-13} , $NR^{VII-7}R^{VII-8}$, halo, $(C=O)C_{1-7}alkyl$, or $SO_m^{VII}R^{VII-9}$, or
- q) R^{VII-1} together with R^{VII-2} form a carbocyclic or het^{VII} which may be optionally substituted by $NR^{VII-7}R^{VII-8}$, or $C_{1-7}alkyl$ which may be optionally substituted by OR^{VII-14} ;

R^{VII-6} is

- a) $C_{1-7}alkyl$,
- b) $NR^{VII-7}R^{VII-8}$
- c) $aryl^{VII}$, or
- d) het^{VII} , wherein said het^{VII} is bound via a carbon atom;

R^{VII-7} and R^{VII-8} are independently

- a) H,
- b) $aryl^{VII}$,
- c) $C_{1-7}alkyl$ which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VII-10}R^{VII-10}$, R^{VII-11} , SO_mR^{VII-9} , $CONR^{VII-10}R^{VII-10}$, or halo, or,
- d) R^{VII-7} and R^{VII-8} together with the nitrogen to which they are attached form a het^{VII} ;

R^{VII-9} is

- a) $aryl^{VII}$,
- b) het^{VII} ,
- c) $C_{3-8}cycloalkyl$,
- d) methyl, or
- e) $C_{2-7}alkyl$ which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VII-10}R^{VII-10}$, R^{VII-11} , SH, $CONR^{VII-10}R^{VII-10}$, or halo;

R^{VII-10} is

- a) H,
- b) methyl, or
- c) $C_{2-7}alkyl$ optionally substituted by OH;

R^{VII-11} is

- a) OR^{VII-10} ,

- b) Ohet^{VII},
- c) Oaryl^{VII},
- d) CO₂R^{VII-10},
- e) het^{VII},
- f) aryl^{VII},
- g) CN, or
- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VII-11}, NR^{VII-7}R^{VII-8}, SO_m^{VII}R^{VII-9}, or C₁₋₇alkyl optionally substituted by R^{VII-11}, NR^{VII-7}R^{VII-8}, or SO_mR^{VII-9};

R^{VII-12} is

- a) H,
- b) het^{VII},
- c) aryl^{VII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8} or R^{VII-11};

R^{VII-13} is

- a) (P=O)(OR^{VII-14})₂,
- b) CO(CH₂)_nCON(CH₃) - (CH₂)_nSO₃⁻M⁺,
- c) an amino acid,
- d) C(=O)aryl^{VII}, or
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8}, aryl^{VII}, het^{VII}, CO₂H, or O(CH₂)_n^{VII}CO₂R^{VII-14};

R^{VII-14} is

- a) H, or
- b) C₁₋₇alkyl;

each n^{VII} is independently 1, 2, 3, 4 or 5;

each m^{VII} is independently 0, 1, or 2;

M^{VII} is sodium, potassium, or lithium;

aryl^{VII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;


wherein any aryl^{VII} is optionally substituted with one or

more substituents selected from the group consisting of halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VII}-14}$, CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VII}-14}$, $\text{NR}^{\text{VII}-14}\text{R}^{\text{VII}-14}$, $\text{OR}^{\text{VII}-14}$, or $\text{CO}_2\text{R}^{\text{VII}-14}$ groups;

het^{VII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VII}-14}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VII}-14}$, $\text{NR}^{\text{VII}-14}\text{R}^{\text{VII}-14}$, $\text{OR}^{\text{VII}-14}$, or $\text{CO}_2\text{R}^{\text{VII}-14}$ groups;

wherein Formula VIII is



and pharmaceutically acceptable salts thereof,

wherein

A^{VIII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO_2 , or
- e) F;

$\text{R}^{\text{VIII}-1}$ is

- a) $\text{R}^{\text{VIII}-5}$,
- b) $\text{NR}^{\text{VIII}-7}\text{R}^{\text{VIII}-8}$, or
- c) $\text{SO}_2\text{R}^{\text{VIII}-9}$;

R^{VIII-2} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) $SO_m R^{VIII-6}$,
- d) OC_{2-7} alkyl substituted by OH,
- e) SC_{2-7} alkyl substituted by OH, or
- f) C_{2-8} alkyl which is partially unsaturated and is optionally substituted by one or more substituents selected from $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl or $SO_m R^{VIII-9}$;

with the proviso that when $R^{VIII-1} = R^{VIII-5} = (CH_2CH_2O)_i R^{VIII-10}$, then R^{VIII-2} may additionally represent

- a) H,
- b) halo,
- c) $(C=O)R^{VIII-6}$,
- d) $(C=O)OR^{VIII-9}$,
- e) cyano,
- f) $OR^{VIII-10}$,
- g) het^{VIII},
- h) $NR^{VIII-7}R^{VIII-8}$,
- i) $SR^{VIII-10}$,
- j) het^{VIII},
- k) $NHCOR^{VIII-12}$,
- l) $NHSO_2R^{VIII-12}$, or
- m) R^{VIII-2} together with R^{VIII-3} or R^{VIII-4} form a carbocyclic or het^{VIII} which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-3} and R^{VIII-4} are independently:

- a) H,
- b) halo,
- c) aryl^{VIII},
- d) $S(O)_m R^{VIII-6}$,
- e) $(C=O)R^{VIII-6}$,

- f) $(C=O)OR^{VIII-9}$,
- g) cyano,
- h) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- i) $OR^{VIII-10}$,
- j) $Ohet^{VIII}$,
- k) $NR^{VIII-7}R^{VIII-8}$,
- l) $SR^{VIII-10}$,
- m) $Shet^{VIII}$,
- n) $NHCOR^{VIII-12}$,
- o) $NHSO_2R^{VIII-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-10}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VIII}RV^{VIII-9}$, or
- q) R^{VIII-4} together with R^{VIII-3} form a carbocyclic or het which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-5} is

- a) $(CH_2CH_2O)_iR^{VIII-10}$,
- b) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- c) aryl VIII ,
- d) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-7}R^{VIII-8}$, $R^{VIII-11}$, SO_mR^{VIII-9} , or OC_{2-4} alkyl which may be further substituted by het^{VIII} , $OR^{VIII-10}$, or $NR^{VIII-7}R^{VIII-8}$, or
- e) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$, $SO_m^{VIII}R^{VIII-9}$, or C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-10}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VIII}RV^{VIII-9}$, or

7alkyl optionally substituted by $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$,
or $SO_m^{VIII}R^{VIII-9}$;

R^{VIII-6} is

- a) C_{1-7} alkyl,
- b) $NR^{VIII-7}R^{VIII-8}$,
- c) aryl^{VIII}, or
- d) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom;

R^{VIII-7} and R^{VIII-8} are independently

- a) H,
- b) aryl^{VIII},
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-10}R^{VIII-10}$, $R^{VIII-11}$, $SO_m^{VIII}R^{VIII-9}$, $CONR^{VIII-10}R^{VIII-10}$, or halo, or,
- d) R^{VIII-7} and R^{VIII-8} together with the nitrogen to which they are attached form a het^{VIII};

R^{VIII-9} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) C_{3-8} cycloalkyl,
- d) methyl, or
- e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-10}R^{VIII-10}$, $R^{VIII-11}$, SH, $CONR^{VIII-10}R^{VIII-10}$, or halo;

$R^{VIII-10}$ is

- a) H,
- b) methyl, or
- c) C_{2-7} alkyl optionally substituted by OH;

$R^{VIII-11}$ is

- a) $OR^{VIII-10}$,

- b) Ohet^{VIII},
- c) Oaryl^{VIII},
- d) CO₂R^{VIII-10},
- e) het^{VIII},
- f) aryl^{VIII}, or
- g) CN;

R^{VIII-12} is

- a) H,
- b) het^{VIII},
- c) aryl^{VIII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VIII-7}R^{VIII-8} or R^{VIII-11};

R^{VIII-13} is

- a) (P=O)(OR¹⁴)₂,
- b) CO(CH₂)_n^{VIII}CON(CH₃)-(CH₂)_n^{VIII}SO₃⁻M⁺,
- c) an amino acid,
- d) C(=O)aryl^{VIII}, or
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VIII-7}R^{VIII-8}, aryl^{VIII}, het^{VIII}, CO₂H, or O(CH₂)_n^{VIII}CO₂R^{VIII-14};

R^{VIII-14} is

- a) H, or
- b) C₁₋₇alkyl;

each i^{VIII} is independently 2, 3, or 4;

each n^{VIII} is independently 1, 2, 3, 4 or 5;

each m^{VIII} is independently 0, 1, or 2;

M^{VIII} is sodium, potassium, or lithium;

aryl^{VIII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;


wherein any aryl^{VIII} is optionally substituted with one

or more substituents selected from halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VIII}-14}$, CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VIII}-14}$, $\text{NR}^{\text{VIII}-14}\text{R}^{\text{VIII}-14}$, $\text{OR}^{\text{VIII}-14}$, or $\text{CO}_2\text{R}^{\text{VIII}-14}$ groups;

het^{VIII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VIII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VIII}-14}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VIII}-14}$, $\text{NR}^{\text{VIII}-14}\text{R}^{\text{VIII}-14}$, $\text{OR}^{\text{VIII}-14}$, or $\text{CO}_2\text{R}^{\text{VIII}-14}$ groups;

wherein Formula IX is



IX

and pharmaceutically acceptable salts thereof, wherein,

$\text{R}^{\text{IX}-1}$ is

- a) Cl,
- b) Br,
- c) CN,
- d) NO_2 , or
- e) F;

$\text{R}^{\text{IX}-2}$, $\text{R}^{\text{IX}-3}$ and $\text{R}^{\text{IX}-4}$ are independently selected from:

- a) H,
- b) halo,
- c) aryl^{IX},

- d) $S(O)_m R^{IX-6}$,
- e) $(C=O) R^{IX-6}$,
- f) $(C=O) OR^{IX-9}$,
- g) cyano,
- h) het^{IX} , wherein said ^{IX}het is bound via a carbon atom,
- i) OR^{IX-10} ,
- j) $Ohet^{IX}$,
- k) $NR^{IX-7}R^{IX-8}$
- l) SR^{IX-10} ,
- m) $Shet^{IX}$,
- n) $NHCOR^{IX-12}$,
- o) $NHSO_2R^{IX-12}$, or
- p) $C_{1-7}alkyl$ which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{IX-11} , OR^{IX-13} , SR^{IX-10} , SR^{IX-13} , $NR^{IX-7}R^{IX-8}$, halo, $(C=O)C_{1-7}alkyl$, or $SO_m R^{IX-9}$;

R^{IX-6} is

- a) $C_{1-7}alkyl$,
- b) $NR^{IX-7}R^{IX-8}$,
- c) $aryl^{IX}$, or
- d) het^{IX} , wherein said het^{IX} is bound via a carbon atom;

R^{IX-7} and R^{IX-8} are independently

- a) H,
- b) $aryl^{IX}$,
- c) $C_{1-7}alkyl$ which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{IX-10}R^{IX-10}$, R^{IX-11} , $SO_m R^{IX-9}$, $CONR^{IX-10}R^{IX-10}$, or halo, or,
- d) R^{IX-7} and R^{IX-8} together with the nitrogen to which they are attached form a ^{IX}het ;

R^{IX-9} is

- a) $aryl^{IX}$,
- b) het^{IX} ,

- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SH, CONR^{IX-10}R^{IX-10}, or halo;

R^{IX-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{IX-11} is

- a) OR^{IX-10},
- b) Ohet^{IX},
- c) Oaryl^{IX},
- d) CO₂R^{IX-10},
- e) het^{IX},
- f) aryl^{IX}, or
- g) CN;

R^{IX-12} is

- a) H,
- b) het^{IX},
- c) aryl^{IX},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{IX-7}R^{IX-8} or R^{IX-11};

R^{IX-13} is

- a) (P=O) (OR^{IX-14})₂,
- b) CO(CH₂)_n^{IX}CON(CH₃) - (CH₂)_n^{IX}SO₃⁻M^{IX+},
- c) an amino acid,
- d) C(=O)aryl^{IX}, or
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{IX-7}R^{IX-8}, aryl^{IX}, het^{IX}, CO₂H, or O(CH₂)_nCO₂R^{IX-14};

R^{IX-14} is

- a) H, or
- b) C_{1-7} alkyl;

each n^{IX} is independently 1, 2, 3, 4 or 5;

each m^{IX} is independently 0, 1, or 2;

M^{IX} is sodium, potassium, or lithium;


aryl^{IX} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three SR^{IX-14} , $NR^{IX-14}R^{IX-14}$, OR^{IX-14} , or CO_2R^{IX-14} groups;

het^{IX} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three SR^{IX-14} , $NR^{IX-14}R^{IX-14}$, OR^{IX-14} , or CO_2R^{IX-14} groups.

2. (Original) The method of claim 1, wherein the compound administered has the Formula



or a pharmaceutically acceptable salt thereof,
wherein,

A^{VI} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VI-1} is

- a) R^{VI-5} , or
- b) SO₂ R^{VI-9}

R^{VI-2} , R^{VI-3} and R^{VI-4} may be the same or different and are selected from the group consisting of:

- a) H,
- b) halo,
- c) aryl^{VI},
- d) S(O)_m R^{VI-6} ,
- e) (C=O) R^{VI-6} ,
- f) (C=O)OR^{VI-9},
- g) cyano,
- h) het^{VI}, wherein said het^{VI} is bound via a carbon atom,
- i) OR^{VI-10},
- j) Ohet^{VI},
- k) NR^{VI-7} R^{VI-8}
- l) SR^{VI-10},
- m) Shet^{VI},
- n) NHCOR^{VI-12},
- o) NHSO₂ R^{VI-12} ,

- p) C₁₋₇alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{VI-11}, OR^{VI-13}, SR^{VI-10}, SR^{VI-13}, NR^{VI-7}R^{VI-8}, halo, (C=O)C₁₋₇alkyl, or SO_m^{VI}R^{VI-9}, and
- q) R^{VI-3} together with R^{VI-2} or R^{VI-4} form a carbocyclic or het^{VI} which may be optionally substituted by NR^{VI-7}R^{VI-8}, or C₁₋₇alkyl which may be optionally substituted by OR^{VI-14};

R^{VI-5} is

- a) (CH₂CH₂O)_i^{VI}R^{VI-10},
- b) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of NR^{VI-7}R^{VI-8}, R^{VI-11}, SO_m^{VI}R^{VI-9}, or OC₂₋₄alkyl which may be further substituted by het^{VI}, OR^{VI-10}, or NR^{VI-7}R^{VI-8}, or
- c) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_m^{VI}R^{VI-9}, or C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}R^{VI-8}, or SO_m^{VI}R⁹;

R^{VI-6} is

- a) C₁₋₇alkyl,
- b) NR^{VI-7}R^{VI-8},
- c) aryl^{VI}, or
- d) het^{VI}, wherein said het^{VI} is bound via a carbon atom;

R^{VI-7} and R^{VI-8} are independently

- a) H,
- b) aryl^{VI},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of aryl^{VI}, NR^{VI-10}R^{VI-10}, R^{VI-11}, SO_m^{VI}R^{VI-9}, CONR^{VI-10}R^{VI-10}, or halo, or;
- d) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents

selected from a group consisting of R^{VI-11} , $NR^{VI-7}R^{VI-8}$, $SO_m^{VI}R^{VI-9}$, or

C_{1-7} alkyl optionally substituted by R^{VI-11} , $NR^{VI-7}R^{VI-8}$, or $SO_m^{VI}R^{VI-9}$, or

- e) R^{VI-7} and R^{VI-8} together with the nitrogen to which they are attached form a het^{VI} ;

R^{VI-9} is

- a) $aryl^{VI}$,
- b) het^{VI} ,
- c) C_{3-8} cycloalkyl,
- d) methyl, or
- e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $NR^{VI-10}R^{VI-10}$, R^{VI-11} , SH, $CONR^{VI-10}R^{VI-10}$, or halo;

R^{VI-10} is

- a) H,
- b) methyl, or
- c) C_{2-7} alkyl optionally substituted by OH;

R^{VI-11} is

- a) OR^{10} ,
- b) $Ohet^{VI}$,
- c) $Oaryl^{VI}$,
- d) CO_2R^{10} ,
- e) het^{VI} ,
- f) $aryl^{VI}$,
- g) CN, or
- h) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11} , $NR^{VI-7}R^{VI-8}$, $SO_m^{VI}R^{VI-9}$, or C_{1-7} alkyl optionally substituted by R^{VI-11} , $NR^{VI-7}R^{VI-8}$, or $SO_m^{VI}R^{VI-9}$;

R^{VI-12} is

- a) H,
- b) het^{VI} ,

- c) aryl^{VI},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VI-7}R^{VI-8} or R^{VI-11};

R^{VI-13} is

- a) (P=O) (OR^{VI-14})₂,
- b) CO(CH₂)_n^{VI}CON(CH₃)-(CH₂)_nSO₃⁻M^{VI+},
- c) an amino acid,
- d) C(=O)aryl^{VI},
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VI-7}R^{VI-8}, aryl^{VI}, het^{VI}, CO₂H, or O(CH₂)_n^{VI}CO₂R^{VI-14}, or
- f) C(=O)NR^{VI-7}R^{VI-8}

R^{VI-14} is

- a) H, or
- b) C₁₋₇alkyl;

each i^{VI} is independently 2, 3, or 4;

each n^{VI} is independently 1, 2, 3, 4 or 5;

each m^{VI} is independently 0, 1, or 2;

M^{VI} is sodium, potassium, or lithium;

aryl^{VI} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO₂R^{VI-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆ alkyl which maybe further substituted by one to three SR^{VI-14}, NR^{VI-14}R^{VI-14}, OR^{VI-14}, or CO₂R^{VI-14};

het^{VI} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VI} is optionally substituted with one or

more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VI}-14}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which maybe further substituted by one to three $\text{SR}^{\text{VI}-14}$, $\text{NR}^{\text{VI}-14}\text{R}^{\text{VI}-14}$, $\text{OR}^{\text{VI}-14}$, or $\text{CO}_2\text{R}^{\text{VI}-14}$.

3. (Original) The method of Claim 2, wherein A^{VI} is Cl.

4. (Original) The method of Claim 2, wherein the compound administered is selected from the group consisting of N-(4-chlorobenzyl)-6-iodo-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-([(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-([(1R,2S)-2-hydroxycyclopentyl]ethynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(2-hydroxyethoxy)methyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-3-furanyl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-(1,2-diethyl-4-pyrazolidinyl)-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-1-(3-oxetanyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-{3-[(3-hydroxypropyl)sulfonyl]propyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[2-(2-ethoxyethoxy)ethyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfinyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-
[(phenylsulfonyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-
[(phenylsulfanyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-
2H-pyran-3-yl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(methylsulfanyl)methyl]-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[[(4-chlorophenyl) sulfinyl]methyl]-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-
2H-pyran-4-yl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-
(4-thiomorpholinylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(4-hydroxy-1-piperidinyl)methyl]-1-
methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[(3R)-3-hydroxypyrrolidinyl]methyl]-1-
methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-hydroxy-1-piperidinyl)methyl]-1-
methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

[3-{{(4-chlorobenzyl)amino}carbonyl}-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinyl]methyl 4-morpholinecarboxylate;

N-(4-chlorobenzyl)-8-(hydroxymethyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-cyanobenzyl)amino]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6,8-bis(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-[(1-acetyl-4-piperidinyl)amino]-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[[1-methyl-2-(phenylsulfonyl)ethyl]amino]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[3-(4-methoxyphenyl)-1-methylpropyl]amino]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-amino-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-8-[(3-nitrobenzyl)amino]-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-(tetrahydro-2H-pyran-4-ylamino)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethyl}-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[(1R,2S)-2-hydroxycyclopentyl]ethyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;


N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-{[methyl(tetrahydro-2-furanylmethyl)amino]methyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

5. (Original) The method of Claim 1, wherein the compound administered has the Formula VII



or a pharmaceutically acceptable salt thereof, wherein,

A^{VII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VII-1} is

- a) aryl^{VII},
- b) S(O)_m^{VII}R^{VII-6},
- c) (C=O)R^{VII-6}, with the proviso that if R^{VII-6} is NR^{VII-7}R^{VII-8}, then R^{VII-7} and R^{VII-8} do not both equal H
- d) (C=O)OR^{VII-9},
- e) cyano,
- f) het^{VII}, wherein said het^{VII} is bound via a carbon atom,
- g) Ohet^{VII},
- h) NR^{VII-7}R^{VII-8} with the proviso that R^{VII-7} and R^{VII-8} do not both equal H
- i) SR^{VII-10},
- j) Shet^{VII},

- k) $\text{NHCOR}^{\text{VII-12}}$,
- l) $\text{NH}\text{SO}_2\text{R}^{\text{VII-12}}$,
- m) $\text{C}_{1-7}\text{alkyl}$ which is partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}\text{alkyl}$, or $\text{SO}_m\text{R}^{\text{VII-9}}$, or
- n) $\text{C}_{1-7}\text{alkyl}$ which is substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}\text{alkyl}$, or $\text{SO}_m\text{R}^{\text{VII-9}}$;

$\text{R}^{\text{VII-2}}$ is

- a) H,
- b) halo,
- c) aryl^{VII} ,
- d) $\text{S}(\text{O})_m\text{R}^{\text{VII-6}}$,
- e) $(\text{C}=\text{O})\text{R}^{\text{VII-6}}$,
- f) $(\text{C}=\text{O})\text{OR}^{\text{VII-9}}$,
- g) cyano,
- h) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- i) $\text{OR}^{\text{VII-10}}$,
- j) Ohet^{VII} ,
- k) $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$,
- l) $\text{SR}^{\text{VII-10}}$,
- m) Shet^{VII} ,
- n) $\text{NHCOR}^{\text{VII-12}}$,
- o) $\text{NH}\text{SO}_2\text{R}^{\text{VII-12}}$, or
- p) $\text{C}_{1-7}\text{alkyl}$ which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}\text{alkyl}$, or $\text{SO}_m\text{R}^{\text{VII-9}}$, or
- q) $\text{R}^{\text{VII-1}}$ together with $\text{R}^{\text{VII-2}}$ form a carbocyclic or het^{VII} which may be optionally substituted by $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, or $\text{C}_{1-7}\text{alkyl}$ which may be optionally substituted by $\text{OR}^{\text{VII-14}}$;

R^{VII-6} is

- a) C₁₋₇alkyl,
- b) NR^{VII-7}R^{VII-8}
- c) aryl^{VII}, or
- d) het^{VII}, wherein said het^{VII} is bound via a carbon atom;

R^{VII-7} and R^{VII-8} are independently

- a) H,
- b) aryl^{VII},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VII-10}R^{VII-10}, R^{VII-11}, SO_mR^{VII-9}, CONR^{VII-10}R^{VII-10}, or halo, or,
- d) R^{VII-7} and R^{VII-8} together with the nitrogen to which they are attached form a het^{VII};

R^{VII-9} is

- a) aryl^{VII},
- b) het^{VII},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VII-10}R^{VII-10}, R^{VII-11}, SH, CONR^{VII-10}R^{VII-10}, or halo;

R^{VII-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VII-11} is

- a) OR^{VII-10},
- b) Ohet^{VII},
- c) Oaryl^{VII},
- d) CO₂R^{VII-10},
- e) het^{VII},
- f) aryl^{VII},
- g) CN, or

- h) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VII-11} , $NR^{VII-7}R^{VII-8}$, $SO_m^{VII}R^{VII-9}$, or C_{1-7} alkyl optionally substituted by R^{VII-11} , $NR^{VII-7}R^{VII-8}$, or $SO_m^{VII}R^{VII-9}$;

R^{VII-12} is

- a) H,
- b) het^{VII} ,
- c) $aryl^{VII}$,
- d) C_{3-8} cycloalkyl,
- e) methyl, or
- f) C_{2-7} alkyl optionally substituted by $NR^{VII-7}R^{VII-8}$ or R^{VII-11} ;

R^{VII-13} is

- a) $(P=O)(OR^{VII-14})_2$,
- b) $CO(CH_2)_n^{VII}CON(CH_3)-(CH_2)_nSO_3^-M^{VII+}$,
- c) an amino acid,
- d) $C(=O)aryl^{VII}$, or
- e) $C(=O)C_{1-7}$ alkyl optionally substituted by $NR^{VII-7}R^{VII-8}$, $aryl^{VII}$, het^{VII} , CO_2H , or $O(CH_2)_n^{VII}CO_2R^{VII-14}$;

R^{VII-14} is

- a) H, or
- b) C_{1-7} alkyl;

each n^{VII} is independently 1, 2, 3, 4 or 5;

each m^{VII} is independently 0, 1, or 2;

M^{VII} is sodium, potassium, or lithium;

$aryl^{VII}$ is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any $aryl^{VII}$ is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO_2R^{VII-14} , CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three SR^{VII-14} , $NR^{VII-14}R^{VII-14}$, OR^{VII-14} , or CO_2R^{VII-14} groups;

het^{VII} is a four- (4), five- (5), six- (6), or seven- (7)

membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VII-14}}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VII-14}}$, $\text{NR}^{\text{VII-14}}\text{R}^{\text{VII-14}}$, $\text{OR}^{\text{VII-14}}$, or $\text{CO}_2\text{R}^{\text{VII-14}}$ groups.

6. (Original) The method of Claim 5, wherein A^{VII} is Cl.

7. (Original) The method of Claim 6, wherein $\text{R}^{\text{VII-1}}$ is selected from the group consisting of CH_2 -morpholine, alkynyl- CH_2OH , CH_2 -(tetrahydro-2H-pyran-4-yl) and $(\text{CH}_2)_3\text{OH}$.

8. (Original) The compound of Claim 6, wherein the compound administered is selected from the group consisting of

N-(4-chlorobenzyl)-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

Methyl 3-{[(4-chlorobenzyl)amino]carbonyl}-4-hydroxy-6-cinnolinecarboxylate;

N-(4-chlorobenzyl)-4-hydroxy-6-(hydroxymethyl)-3-cinnolinecarboxamide N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{[(1R,2S)-2-hydroxycyclopentyl]ethynyl}-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[(1R,2S)-2-hydroxycyclopentyl]ethyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;


N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethyl]-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

9. (Original) A method of Claim 1, wherein the compound administered is Formula VIII



and pharmaceutically acceptable salts thereof, wherein A^{VIII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VIII-1} is

- a) R^{VIII-5} ,
- b) $NR^{VIII-7}R^{VIII-8}$, or
- c) SO_2R^{VIII-9} ;

R^{VIII-2} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) SO_m^{VIII}R^{VIII-6},
- d) OC₂₋₇ alkyl substituted by OH,
- e) SC₂₋₇ alkyl substituted by OH, or
- f) C₂₋₈ alkyl which is partially unsaturated and is optionally substituted by one or more substituents selected from R^{VIII-11}, OR^{VIII-13}, SR^{VIII-13}, NR^{VIII-7}R^{VIII-8}, halo, (C=O)C₁₋₇ alkyl or SO_m^{VIII}R^{VIII-9};

with the proviso that when R^{VIII-1} = R^{VIII-5} = (CH₂CH₂O)_iR^{VIII-10}, then R^{VIII-2} may additionally represent

- a) H,
- b) halo,
- c) (C=O)R^{VIII-6},
- d) (C=O)OR^{VIII-9},
- e) cyano,
- f) OR^{VIII-10},
- g) Ohet^{VIII},
- h) NR^{VIII-7}R^{VIII-8},
- i) SR^{VIII-10},
- j) Shet^{VIII},
- k) NHCOR^{VIII-12},
- l) NHSO₂R^{VIII-12}, or
- m) R^{VIII-2} together with R^{VIII-3} or R^{VIII-4} form a carbocyclic or het^{VIII} which may be optionally substituted by NR^{VIII-7}R^{VIII-8}, or C₁₋₇alkyl which may be optionally substituted by OR^{VIII-14};

R^{VIII-3} and R^{VIII-4} are independently:

- a) H,
- b) halo,
- c) aryl^{VIII},
- d) S(O)_m^{VIII}R^{VIII-6},
- e) (C=O)R^{VIII-6},

- f) $(C=O)OR^{VIII-9}$,
- g) cyano,
- h) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- i) $OR^{VIII-10}$,
- j) $Ohet^{VIII}$,
- k) $R^{VIII-7}R^{VIII-8}$,
- l) $SR^{VIII-10}$,
- m) $Shet^{VIII}$,
- n) $NHCOR^{VIII-12}$,
- o) $NHSO_2R^{VIII-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-10}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VIII}R^{VIII-9}$, or
- q) R^{VIII-4} together with R^{VIII-3} form a carbocyclic or het^{VIII} which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-5} is

- a) $(CH_2CH_2O)_iR^{VIII-10}$,
- b) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- c) $aryl^{VIII}$,
- d) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-7}R^{VIII-8}$, $R^{VIII-11}$, $SO_m^{VIII}R^{VIII-9}$, or OC_{2-4} alkyl which may be further substituted by het^{VIII} , $OR^{VIII-10}$, or $NR^{VIII-7}R^{VIII-8}$, or
- e) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$, $SO_m^{VIII}R^{VIII-9}$, or C_{1-}

7alkyl optionally substituted by $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$,
or SO_mR^{VIII-9} ;

R^{VIII-6} is

- a) C_{1-7} alkyl,
- b) $NR^{VIII-7}R^{VIII-8}$,
- c) aryl^{VIII}, or
- d) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom;

R^{VIII-7} and R^{VIII-8} are independently

- a) H,
- b) aryl^{VIII},
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-10}R^{VIII-10}$, $R^{VIII-11}$, SO_mR^{VIII-9} , $CONR^{VIII-10}R^{VIII-10}$, or halo, or,
- d) R^{VIII-7} and R^{VIII-8} together with the nitrogen to which they are attached form a het^{VIII};

R^{VIII-9} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) C_{3-8} cycloalkyl,
- d) methyl, or
- e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-10}R^{VIII-10}$, $R^{VIII-11}$, SH, $CONR^{VIII-10}R^{VIII-10}$, or halo;

$R^{VIII-10}$ is

- a) H,
- b) methyl, or
- c) C_{2-7} alkyl optionally substituted by OH;

$R^{VIII-11}$ is

- a) $OR^{VIII-10}$,
- b) Ohet^{VIII},

- c) Oaryl^{VIII},
- d) CO₂R^{VIII-10},
- e) het^{VIII},
- f) aryl^{VIII}, or
- g) CN;

R^{VIII-12} is

- a) H,
- b) het^{VIII},
- c) aryl^{VIII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VIII-7}R^{VIII-8} or R^{VIII-11};

R^{VIII-13} is

- a) (P=O)(OR¹⁴)₂,
- b) CO(CH₂)_n^{VIII}CON(CH₃)-(CH₂)_nSO₃⁻M^{VIII+},
- c) an amino acid,
- d) C(=O)aryl^{VIII}, or
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VIII-7}R^{VIII-8}, aryl^{VIII}, het^{VIII}, CO₂H, or O(CH₂)_n^{VIII}CO₂R^{VIII-14};

R^{VIII-14} is

- a) H, or
- b) C₁₋₇alkyl;

each i^{VIII} is independently 2, 3, or 4;

each n^{VIII} is independently 1, 2, 3, 4 or 5;

each m^{VIII} is independently 0, 1, or 2;

M^{VIII} is sodium, potassium, or lithium;

aryl^{VIII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic; wherein any aryl^{VIII} is optionally substituted with one or more substituents selected from halo, OH, cyano, CO₂R^{VIII-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆ alkyl which may be

further substituted by one to three $\text{SR}^{\text{VIII-14}}$, $\text{NR}^{\text{VIII-14}}\text{R}^{\text{VIII-14}}$, $\text{OR}^{\text{VIII-14}}$, or $\text{CO}_2\text{R}^{\text{VIII-14}}$ groups; het^{VIII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group; wherein any het^{VIII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VIII-14}}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VIII-14}}$, $\text{NR}^{\text{VIII-14}}\text{R}^{\text{VIII-14}}$, $\text{OR}^{\text{VIII-14}}$, or $\text{CO}_2\text{R}^{\text{VIII-14}}$ groups.

10. (Original) The method of Claim 9, wherein A^{VIII} is Cl.

11. (Original) The method of Claim 9, wherein $\text{R}^{\text{VIII-2}}$ is alkynyl- CH_2OH .

12. (Original) The method of Claim 9, wherein the compound administered is N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide, or N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide; or a pharmaceutically acceptable salt thereof.

13. (Original) The method of Claim 9, wherein the compound administered is:

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-Chlorobenzyl)-6-iodo-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1,7-dimethyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1-methyl-4,7-dioxo-1,4,7,8-tetrahydro[1,8]naphthyridine-3-carboxamide;


N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

ethyl 6-([(4-chlorobenzyl)amino]carbonyl)-2-methoxy-8-methyl-5-oxo-5,8-dihydro[1,8]naphthyridine-3-carboxylate;

and pharmaceutically acceptable salts thereof.

14. (Original) A method of Claim 1, wherein the compound administered has the Formula IX



IX

and pharmaceutically acceptable salts thereof, wherein,

R^{IX-1} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO_2 , or
- e) F;

R^{IX-2} , R^{IX-3} and R^{IX-4} are independently selected from:

- a) H,
- b) halo,
- c) $aryl^{IX}$,
- d) $S(O)_m R^{IX-6}$,
- e) $(C=O) R^{IX-6}$,
- f) $(C=O) OR^{IX-9}$,
- g) cyano,
- h) het^{IX} , wherein said het^{IX} is bound via a carbon atom,
- i) OR^{IX-10} ,
- j) $Ohet^{IX}$,
- k) $NR^{IX-7} R^{IX-8}$
- l) SR^{IX-10} ,
- m) $S^{IX-}het$,
- n) $NHCO R^{IX-12}$,
- o) $NHSO_2 R^{IX-12}$, or
- p) $C_{1-7}alkyl$ which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{IX-11} , OR^{IX-13} , SR^{IX-10} , SR^{IX-13} , $NR^{IX-7} R^{IX-8}$, halo, $(C=O)C_{1-7}alkyl$, or $SO_m R^{IX-9}$;

R^{IX-6} is

- a) $C_{1-7}alkyl$,
- b) $NR^{IX-7} R^{IX-8}$,
- c) $aryl^{IX}$, or
- d) het^{IX} , wherein said het^{IX} is bound via a carbon atom;

R^{IX-7} and R^{IX-8} are independently

- a) H,

- b) aryl^{IX},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SO_mR^{IX-9}, CONR^{IX-10}R^{IX-10}, or halo, or,
- d) R^{IX-7} and R^{IX-8} together with the nitrogen to which they are attached form a het^{IX};

R^{IX-9} is

- a) aryl^{IX},
- b) het^{IX},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SH, CONR^{IX-10}R^{IX-10}, or halo;

R^{IX-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{IX-11} is

- a) OR^{IX-10},
- b) Ohet^{IX},
- c) Oaryl^{IX},
- d) CO₂R^{IX-10},
- e) het^{IX},
- f) aryl^{IX}, or
- g) CN;

R^{IX-12} is

- a) H,
- b) het^{IX},
- c) aryl^{IX},
- d) C₃₋₈cycloalkyl,

- e) methyl, or
 - f) C_{2-7} alkyl optionally substituted by $NR^{IX-7}R^{IX-8}$ or R^{IX-11} ;
- R^{IX-13} is
- a) $(P=O)(OR^{IX-14})_2$,
 - b) $CO(CH_2)_n^{IX}CON(CH_3)-(CH_2)_n^{IX}SO_3^-M^{IX+}$,
 - c) an amino acid,
 - d) $C(=O)$ aryl, or
 - e) $C(=O)C_{1-7}$ alkyl optionally substituted by $NR^{IX-7}R^{IX-8}$, aryl^{IX}, het^{IX}, CO_2H , or $O(CH_2)_nCO_2R^{IX-14}$;

R^{IX-14} is

- a) H, or
- b) C_{1-7} alkyl;

each n^{IX} is independently 1, 2, 3, 4 or 5;

each m^{IX} is independently 0, 1, or 2;

M^{IX} is sodium, potassium, or lithium;

aryl^{IX} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three SR^{IX-14} , $NR^{IX-14}R^{IX-14}$, OR^{IX-14} , or CO_2R^{IX-14} groups;

het^{IX} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, oxo,

oxime, and C₁₋₆ alkyl which may be further substituted by one to three SR^{IX-14}, NR^{IX-14}R^{IX-14}, OR^{IX-14}, or CO₂R^{IX-14} groups.

15. (Original) The method of Claim 14, wherein R^{IX-1} is Cl.

16. (Original) The method of Claim 14, wherein the compound administered is selected from a group consisting of

N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(tetrahydro-2H-pyran-4-ylmethyl)[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(4-morpholinylmethyl)[1,8]naphthyridine-3-carboxamide;

6-bromo-N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-iodo-7-methyl[1,8]naphthyridine-3-carboxamide; and

Methyl 6-{[(4-chlorobenzyl)amino]carbonyl}-5-hydroxy-2-methyl[1,8]naphthyridine-3-carboxylate.

17. (Original) The method according to Claim 1, wherein said mammal is a human.

18. (Original) The method according to Claim 1, wherein said mammal is a livestock or companion animal.

19. (Original) The method according to Claim 1, wherein the amount administered is from about 0.1 to about 300 mg/kg of mammal body weight.

20. (Original) The method according to Claim 1, wherein the amount administered is from about 1 to about 30 mg/kg of mammal body weight.

21. (Currently Amended) The method according to ~~Claim 2~~Claim 1, wherein the compound is administered parenterally, intravaginally, intranasally, topically, orally, or rectally. |